

I claim:

1. A method for treating a skin disorder, the method comprising a step of administering a botulinum toxin to a location of a skin disorder of a patient, thereby treating the skin disorder.
2. The method of claim 1, wherein the botulinum toxin is selected from the group consisting of botulinum toxin types A, B, C, D, E, F and G.
3. The method of claim 1, wherein the botulinum toxin is a botulinum toxin type A.
4. The method of claim 1, wherein the botulinum toxin is administered in an amount of between about 1 unit and about 3,000 units.
5. The method of claim 1, wherein the administration is by topical or subcutaneous administration of the botulinum toxin.
6. A method for treating a skin disorder, the method comprising the step of locally administering between 1 unit and 3000 units of a botulinum toxin to a skin disorder of the patient, thereby treating the skin disorder.
7. The method of claim 6, wherein the skin disorder is selected from the group consisting of a wart, corn, callus and bunion.
8. The method of claim 6, wherein the skin disorder is treated by reducing a pain associated with the skin disorder.
9. The method of claim 6, wherein the skin disorder is treated by reducing an inflammation associated with the skin disorder.

10. The method of claim 6, wherein the skin disorder is treated by reducing a size of the skin disorder.

11. A method for treating a wart, the method comprises the step of
5 administering an effective amount of a botulinum toxin to a wart, thereby treating the wart.